Schelkovo Agrohim
Zavodskaya street 2
141108 Schelkovo
Russia
+7 495 745 0551, 745 0198
info@betaren.ru

Ultramag Calcium Active

Fluid concentrated fertilizer with a high content of calcium in the highly absorbable plant form for foliar nutrition of agricultural, fruit and vegetable crops. A complex foliar fertilizer for crop nutrition containing amino acids, calcium, and microelements.

Advantages:

— Rapid and effective uptake and adsorption of calcium and boron
— Calcium, boron, and zinc deficiency compensation
— Improves the quality of flowering and fruit setting
— Increases stress resistance due to its amino acid content
— Improves crop yields and quality

Usage regulations

Crop	Method, time, features of application	Working liquid consumption rate, l/ha	Working liquid consumption rate, l/ha
Cereal crops	Foliar application during tillering and from stalk-shooting to flowering	1-2	200-400
Sugar beet, fodder beet, table beet	Foliar application during the 4–6 leaves stage, 20–25 days after the first application, and 14–20 days before harvesting	1-2	200-400
Maize	Foliar application during the 4–6 leaves stage and 10–14 days after the first application	1-2	200-400
Legumes, legume grasses	Foliar application during the budding phase — early flowering, and at the end of flowering – during pod formation	1-2	200-400
Winter rapeseed, spring rapeseed, winter cress, false flax	Foliar application in spring, when the growing period is resumed, or at the phase of fully sprouted plants and budding/flowering phase	1-2	200-400
Sunflower	Foliar application from the start of pseudanthium formation till flowering, 1–2 times	1-2	200-400
Vegetable crops, potato	Foliar application during budding/start of flowering, and then 2–3 times every 10–15 days	1-4	200-400
Fiber flax, oilseed flax	Foliar application during the herringbone phase and budding	1-2	200-400

Fruit crops (pome fruit and stone fruit crops)	Foliar application before flowering, at the end of flowering, and during fruit growth, 1–2 times	1-4	600-1000
Grapes	Foliar application during flower initiation/start of flowering and during berry growth, 2–3 times every 14–21 days	1-4	600-800
Berries	Foliar application during budding and start of flowering	1-4	200-400

General information

Foliar fertilisers Ultramag Calcium (content in %wt)

Content	%
N	2,0
CaO	9,0
Zn	1.4
B	1.2
	3.7

Shelf life

2 years

Registrant

Schelkovo Agrohim, Russia

Manufacturer

Schelkovo Agrohim, Russia