



Ultramix Development

Liquid mineral fertilizer with microelements for foliar feeding of fruit, fruit and berry, vegetable crops, and vineyards on all types of soil.

Action

Composition of fertilisers Ultramix, %

Brands	N total	CaO	MgO	Fe	Mn	Zn	Cu	B	Mo
Development	2,0	10,0	5,0	0,7	2,0	-	-	-	-

Usage regulations

Crop	Product application dose	Method, time and features of application
Fruit crops (pome fruit and stone fruit crops)	1.0-3.0 L/ha Working solution consumption 600-1,000 L/ha	Foliar feeding of plants is recommended to be carried out 1-2 times: the 1st feeding is carried out during the fruit growth phase (40-55 mm), the 2nd feeding takes place after 14-21 days
Grapes	1.0-3.0 L/ha Working solution consumption: 600-1,000 L/ha	Foliar feeding of plants is recommended to be carried out 1-2 times: the 1st feeding is carried out during the bunch formation, the 2nd feeding takes place after 14-21 days
Berry crops	1.0-2.0 L/ha Working solution consumption: 200-600 L/ha	Foliar feeding of plants after flowering and then 1-2 times with an interval of 14-21 days
Vegetable crops	1.0-2.0 L/ha Working solution consumption: 200-600 L/ha	Foliar feeding of plants during the period of fruit formation and growth 1-3 times with an interval of 14-21 days