



Drotik Combi

Drotik Combi

colloidal solution concentrate and wettable powder

2,4-D acid /2-ethylhexyl ether/ 400 g/L+ metsulfuronmethyl 600 g/kg

Binary packs of systemic postemergence herbicides with selective action against a wide range of dicotyledonous weeds in cereal crops.

Advantages:

- Maximum protection at minimum cost
- Convenient packaging with pre-measured doses for preparing tank mixtures
- Labor savings and reduced per-hectare treatment costs
- More efficient use of machinery and equipment
- Elimination of mixing errors in tank mixtures
- Enhanced effectiveness against target species and expanded spectrum of action

Action

A binary pack consists of a canister with two compartments, where the liquid and solid components are kept separate.

One binary pack is designed to treat 8–12 hectares of crops and allows precise dosing of mixture components, ensuring no risk of overdose.

Usage regulations

Crop	Harmful object	Application rate for products in the package	Application rate per package	Method, time, features of application	Waiting time, days (number of applications)
------	----------------	--	------------------------------	---------------------------------------	---

<p>Spring and winter wheat, spring and winter barley</p>	<p>Annual dicotyledonous weeds, including 2,4-D and MCPA resistant weeds, perennial dicotyledonous weeds (thistle and sow thistle, lettuce, vine milkweed, sheepbine, etc.)</p>	<p>Drotik, CSC 5L + Zinger, WP 50g (Drotik, CSC 0.4-0.6 L/ha + Zinger, WP 4-6 g/ha) Spray Volume: 200-300 L/ha.</p>	<p>8-12 ha</p>	<p>Spray crops during the tillering stage (for winter cereals, treat in spring). Target weeds in their early growth stages: annuals at the 2-4 leaf stage, perennials at the rosette stage. The following year after harvesting cereals: sugar beet and vegetable crops must not be sown, Sunflower and buckwheat may be sown only after deep ploughing. Do NOT sow sunflower or buckwheat the following year if either of these conditions is met: Soil pH is above 7.5 or a prolonged drought occurred between herbicide application and sowing of these crops. If replanting is necessary on treated areas, sow spring cereals only.</p>	<p>60(1)</p>
--	--	---	----------------	--	--------------

General information

Chemical class

phenoxy-carboxylates + sulfonyleureas

Storage temperature range

minus 20 °C to plus 30 °C

Shelf life

3 years

Hazard class

Hazard class 2, high danger

Packing

Binary packs: Drotik, CSC 5L + Zinger, WP 50g

Registrant

Schelkovo Agrohim, Russia

Manufacturer

Schelkovo Agrohim, Russia