



Ultramag Phosphorus Super

A liquid mineral fertiliser with macro- and micro-nutrients for the foliar dressing of agricultural crops.

Advantages:

- Eliminates pseudodeficiencies and deficiencies of phosphorus in plants
- Fully supplies plants with easily digestible phosphorus during early development stages
- Activates photosynthesis processes in plants
- Increases plant resistance to weather changes and cold temperatures
- Accelerated crop formation

Usage regulations

Crop	Method, time and features of application	Product consumption rate, l/ha	Working liquid consumption rate, l/ha
Cereal crops	Foliar application at the 3–4 leaf stage, at the tillering stage, shooting and panicle stage	2–6	200-400
Maize	Foliar application at the 2–6 leaf stage, 7–8 leaf stage and in the stem elongation stage (before tasselling)	2–6	200-400
Legumes	Foliar application to plants at the 3–4 ternate leaves stage, at the budding stage and the filling stage	2–6	200-400
Sunflower	Foliar application at the 2–3 leaf-pair stage and at the start of the budding stage	2–6	200-400
Sugar beet	Foliar application is at the 4–6 leaf stage, at the 8–10 leaf stage and at the rows closing stage	2–6	200-400
Spring rapeseed	Foliar application at the 4–8 leaf stage, at the budding stage and at the end of blooming	2–6	200-400
Winter rapeseed	Foliar application at the 4–8 leaf stage (in autumn), at the start of vegetation resumption, at the budding stage and at the end of the blooming stage	2–6	200-400

Potato	Foliar application at the full seedlings stage, at the budding stage and after blooming 1-2 times with the 10-14-day interval	2-6	200-400
Cabbage (all types)	Foliar application 2-3 weeks after the seedling planting and at the start of the loaf (rosette) formation	2-6	400-600
Vegetable crops	Foliar application 2-3 weeks after the seedling planting (emergence of seedlings) and then 1-2 times with a 10-15-day interval.	2-6	400-600
Edible roots, onions, garlic, lettuce	Foliar application 7-10 days of the week after the emergence of seedlings (transplanting of seedlings), then 1-2 times at with interval of 10-15 days	2-6	400-600
Fruit and berry crops, grapes	Foliar application at the beginning of the budding stage and then 3-4 times every 10-15 days (the last dressing is 2-4 weeks before harvesting)	4-6	600-1000

General information

Foliar fertilisers Ultramag Phosphorus Super (content in %wt, g/L)

Content	%	g/L
N	6,4	100,0
P ₂ O ₅	35,0	539,0
MgO	4,0	62,0
Zn	2,5	38,0

Storage conditions

Storage temperature range - minus 5 °C to plus 30 °C

Shelf life

2 years

Registrant

Schelkovo Agrohim, Russia

Manufacturer

Schelkovo Agrohim, Russia